



Township Public Schools

Health Services  
50 Nellis Drive  
Wayne, NJ 07470

Phone: (973) 317-2198

Fax: (973) 317-2159

Dear Parent or Guardian,

An estimated 6 to 12 million infestations of Pediculosis (head lice) occur each year in the United States, most commonly among children ages 3 to 11.

**What are head lice?** Pediculosis (head lice) are tiny, wingless insects that live close to the human scalp. They feed on blood. The eggs, are called nits, are tiny, tear-drop shaped eggs that attach to the hair shaft. Nits often appear yellowish or white, and can look like dandruff but cannot be removed or brushed off. The nymph, or baby louse, is smaller and grow to adult size in one to two weeks. The adult louse is the size of a sesame seed, appears tan to grayish-white. An itchy and inflamed scalp is a common symptom of lice. Although not common, persistent scratching may lead to skin irritation and even infection.

**Who is affected by Pediculosis (head lice)?** Head lice are not related to cleanliness. In fact, head lice often infest people with good hygiene and grooming habits. Head lice are mostly spread by direct head-to-head contact – for example, during play at home or school. Slumber parties, sports activities, or camp. Pediculosis can spread via objects that have been in recent contact with a person with head lice, such as hats, scarves, hair ribbons, combs, brushes, stuffed animals or bedding.

**What to do if an infestation occurs?** Careful inspection of the child's hair and scalp particular attention given to the nape of the neck, crown of the head and above and behind the ears is best way to detect signs of infestation. The nits may be more easily recognized by examining in good light (sunlight) or by shining a light on the child's scalp during the examination. If lice is detected or suspected, consult with your physician for prompt treatment of the child and the environment to prevent re-infestation. Family members or other close contacts should also be examined to control the spread of infestation. Early detection and prompt treatment are essential in attempting to control the spread of lice. Regular examination throughout the school year of your child's hair and scalp will be greatly appreciated. Items that cannot be washed or dry-cleaned may be sealed in plastic bags for two or three weeks. Contact the school nurse who can help you with proper identification, counsel you and monitor the effectiveness of treatment while also maintaining your confidentiality. Children should NOT be treated merely on speculation.

If you have any further questions, please contact your physician or school nurse.

Sincerely,

**The Wayne Township Public School Nurse**

# Head Lice 101

## What You Should Know About Head Lice

## Lice Lessons



### Overview

Head lice are a common community problem. An estimated 6 to 12 million infestations occur each year in the United States, most commonly among children ages 3 to 11 years old.<sup>1</sup> Though a head lice infestation is often spotted in school, it is usually acquired through direct head-to-head contact elsewhere, such as at sleepovers or camp.<sup>2</sup>

Head lice are not dangerous, and they do not transmit disease.<sup>1</sup> Additionally, despite what you might have heard, head lice often infest people with good hygiene and grooming habits.<sup>3,4</sup> Your family, friends, or community may experience head lice. It's important to know some basics, including how to recognize symptoms and what to do if faced with an infestation.

### Fast Facts

- An estimated 6 to 12 million infestations occur each year among US children 3 to 11 years of age<sup>1</sup>
- Head lice do not discriminate, often infesting people with good hygiene.<sup>3,4</sup> They spread mainly through head-to-head contact<sup>1</sup>
- If you or your child exhibits signs of an infestation, it is important to talk to your doctor to learn about treatment options

### What are head lice?

Head lice are tiny, wingless insects that live close to the human scalp. They feed on human blood.<sup>1</sup> When checking for head lice, you may see several forms: the nit, the nymph, and the adult louse.



**Nits** are tiny, teardrop-shaped lice eggs that are often yellowish or white. Nits are also what you call the shells that are left behind once the eggs hatch. Nits are attached to the hair shaft and often found around the nape of the neck or the ears. Nits can look similar to dandruff, but cannot be easily removed or brushed off.<sup>1</sup>



**Nymphs**, or baby lice, are small and grow to adult size in 1 to 2 weeks.<sup>1</sup>



**Adult lice** are the size of a sesame seed and appear tan to grayish-white.<sup>1</sup>

### How are head lice spread?

- Head lice move by crawling and cannot jump or fly<sup>1</sup>
- Head lice are mostly spread by direct head-to-head contact—for example, during play at home or school, sleepovers, sports activities, or camp<sup>1</sup>
- It is possible, but not common, to spread head lice by contact with items that have been in contact with a person with head lice, such as clothing (for example, hats, scarves, or coats) or other personal items (such as combs, brushes, or towels)<sup>1</sup>
- Head lice transmission can occur at home, in the community, or—very infrequently—in school<sup>1,2</sup>

### What are the signs and symptoms of infestation?

Signs and symptoms of infestation include<sup>1</sup>:

- **Tickling** feeling on the scalp or in the hair
- **Itching** (caused by the bites of the louse)
- **Irritability and difficulty sleeping** (lice are more active in the dark)
- **Sores on the head** (caused by scratching, which can sometimes become infected)

Finding a live nymph or adult louse on the scalp or in the hair is an indication of an active infestation. They are most commonly found behind the ears and near the neckline at the back of the head.<sup>1</sup>

# Head Lice 101

## What You Should Know About Head Lice

## Lice Lessons



### What if my child gets head lice?

If you suspect your child might have head lice, it's important to talk to a school nurse, pediatrician, or family physician to get appropriate care. There are a number of available treatments, including new prescription treatment options that are safe and do not require nit combing. Other things to consider in selecting and starting treatment include:

- Follow treatment instructions. Using extra amounts or multiple applications of the same medication is not recommended, unless directed by a healthcare professional<sup>5</sup>
- A 2016 study showed that 48 states now have lice that are genetically predisposed to resistance to commonly used treatments<sup>6</sup>
- There is no scientific evidence that home remedies are effective treatments<sup>7</sup>
- Head lice do not infest the house. However, family bed linens and recently used clothes, hats, and towels should be washed in very hot water and dried on the high setting<sup>5</sup>
- Personal articles, such as combs, brushes, and hair clips, should be soaked in very hot water (at least 130°F) for 5 to 10 minutes if they were exposed to someone with an active head lice infestation<sup>5</sup>
- All household members and other close contacts should be checked, and those with evidence of an active infestation should also be treated at the same time<sup>5</sup>

### Myths and facts about head lice

**Myth:** Only dirty people get head lice.

**Fact:** Personal hygiene and household or school cleanliness are not factors for infestation. In fact, head lice often infest people with good hygiene and grooming habits.<sup>3,4</sup>

**Myth:** Head lice carry diseases.

**Fact:** Head lice do not spread diseases.<sup>1</sup>

**Myth:** Head lice can be spread by sharing hair brushes, hats, clothes, and other personal items.

**Fact:** It is uncommon to spread head lice by contact with clothing or other personal items, such as combs, brushes, or hair accessories, that have been in contact with a person with head lice.<sup>1</sup>

**Myth:** Head lice can jump or fly, and can live anywhere.

**Fact:** Head lice cannot jump or fly, and only move by crawling. It is unlikely to find head lice living on objects like helmets or hats because they have feet that are specifically designed to grasp on to the hair shaft of humans. Additionally, a louse can only live for about a day off the head.<sup>1</sup>

**Myth:** You can use home remedies like mayonnaise to get rid of head lice.

**Fact:** There is no scientific evidence that home remedies are effective treatments.<sup>7</sup> Consult your healthcare provider to discuss appropriate treatment options, including prescription products.

### References

1. Centers for Disease Control and Prevention. Frequently asked questions (FAQs). [http://www.cdc.gov/parasites/lice/head/gen\\_info/faqs.html](http://www.cdc.gov/parasites/lice/head/gen_info/faqs.html). Accessed April 20, 2017.
2. Pontius DJ. Demystifying pediculosis: school nurses taking the lead. *Pediatr Nurs*. 2014;40(5):226-235.
3. Meinking T, Taplin D, Vicaria M. Infestations. In: Schachner LA, Hansen RC, eds. *Pediatric Dermatology*, 4th ed. Mosby Elsevier; 2011:1535-1583.
4. Centers for Disease Control and Prevention. Epidemiology & risk factors. <http://www.cdc.gov/parasites/lice/head/epi.html>. Accessed April 20, 2017.
5. Centers for Disease Control and Prevention. Treatment. <http://www.cdc.gov/parasites/lice/head/treatment.html>. Accessed April 20, 2017.
6. Gellatly KJ, Krim S, Palenchar DJ, et al. Expansion of the knockdown resistance frequency map for human head lice (phthiraptera: pediculidae) in the United States using quantitative sequencing. *J Med Entomol*. 2016;1-7.
7. Centers for Disease Control and Prevention (CDC). Treatment frequently asked questions. [http://www.cdc.gov/parasites/lice/head/gen\\_info/faqs\\_treat.html](http://www.cdc.gov/parasites/lice/head/gen_info/faqs_treat.html). Accessed April 20, 2017.

Lice Lessons educational initiative is made possible through a collaboration with Arbor Pharmaceuticals, LLC.

PP-NP-US-0265

arbor  
PHARMACEUTICALS, LLC

NASN  
National  
Association of  
School Nurses